

Coronavirus disease (COVID-19): Characteristics in children and considerations for dentists providing their care

Abstract

The emergence of the novel virus severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) has caused a global pandemic called coronavirus disease (COVID-19) and has become one of the most significant challenges to the healthcare profession. Dental practices are focal points for cross-infection, and care must be taken to minimise the risk of infection to, from, or between dental care professionals and patients. The COVID-19 epidemiological and clinical characteristics are still being collated but children's symptoms seem to be milder than those that adults experience. It is unknown whether certain groups, for example children with comorbidities, might be at a higher risk of more severe illness. Emerging data on disease spread in children, affected by COVID-19, have not been presented in detail. The purpose of this article was to report current data on the paediatric population affected with COVID-19 and highlight considerations for dentists providing care for children during this pandemic. All members of the dental team have a professional responsibility to keep themselves informed of current guidance and be vigilant in updating themselves as recommendations are changing so quickly.

1 | INTRODUCTION

At the beginning of 2020, the novel virus severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) appeared, causing the coronavirus disease (COVID-19). The emerging virus has resulted in a global pandemic declared a Public Health Emergency of International Concern (PHEIC) by the World Health Organization (WHO) Director-General on the recommendation of the International Health Regulations (2005) Emergency Committee.¹ The case detection rate is changing daily and can be tracked in almost real time.² As of 31 March 2020, 19:50 hours (Central Standard Time), the number of confirmed cases was 857 487 and reported deaths were

42 106 with 169 418 recovered patients.² The first case of a dentist being tested positive for COVID-19 was reported on 23 January 2020 at the Department of Preventive Dentistry in the Wuhan University Dental Hospital. Eventually, the transmission of disease to eight other oral healthcare professionals was identified.³ The characteristics of epidemiological spread and clinical manifestations of COVID-19 in children have not yet been thoroughly elucidated. This article reports current data on the paediatric population affected with COVID-19 and emphasises the importance of following locally, regionally, and nationally relevant safety measures to protect dental care professionals as well as the child patient, whilst providing clinical care for the obviously affected children and those potential carriers of the infection. We emphasise that, in a rapidly changing pandemic landscape, practitioners must actively, regularly seek and use reputable and reliable sources of information on managing child patients that are appropriate for their own region and circumstances.

2 | COVID-19

2.1 | Clinical characteristics of COVID-19 in children

The clinical symptoms of COVID-19 are still being documented and collated, although the majority of affected patients exhibit symptoms including a dry cough which is usually accompanied by fever.⁴ Difficulty in breathing, fatigue, and other less typical symptoms can also occur.^{5,6} Signs and symptoms include different stages as asymptomatic, mild, moderate, severe, and critical.⁷ Children tend to present with similar but milder symptoms to adults. To date, 3092 paediatric cases have been reported to have tested positive, and 1412 children were suspected of having been infected with COVID-19. A survey of 1391 children in China found 171 (12.3%) cases tested positive for SARS-CoV-2.⁸ An analysis of more than 2000 child patients with suspected or confirmed COVID-19 in Hubei, China, found that over 90% presented as asymptomatic or with mild to moderate symptoms.⁹ A summary of paediatric cases reported

with COVID-19 is presented in Table 1.^{7,9-45} These numbers are likely to be under-representative, as there is not universal testing of the whole population for the presence of COVID-19. An overall fatality rate of 1.36%-15% has been reported across all patients with COVID-19.⁴⁶ As of 31 March 2020, seven fatalities have been reported in paediatric population due to COVID-19. These are an infant in Chicago,⁴⁷ a minor child in New York,⁴⁸ a 12-year-old boy in Belgium,⁴⁹ a 13-year-old boy in England,⁵⁰ a 14-year-old boy in China,⁷ a 16-year-old girl in France,⁵¹ and a 17-year-old boy in Los Angeles.⁵²

2.2 | The child patient in the dental setting

Because of the long incubation period (2-14 days)⁶ for everyone, and because children can be asymptomatic or present with mild, nonspecific symptoms, all child patients and parents should be considered as potential carriers of COVID-19 unless proved otherwise. COVID-19 can be transmitted through direct and indirect contact, mainly via respiratory droplets and splatter from saliva and blood through contact with mucous membranes and contaminated fomites.^{53,54} This leaves dental professionals in potentially high-risk situations. Many dental treatments are aerosol generating procedures (AGPs), which have been associated with the transmission of acute respiratory infections.⁵⁵ In addition, dental settings are more likely to have a high number of potentially contaminated surfaces such as dental chairs, their handles, the spittoon, and dental instruments after carrying out a treatment which are possible routes of transmission.³ SARS-CoV-2 virus can persist on surfaces for up to 72 hours,⁵⁶ and all clinic surfaces should be disinfected using chemicals recommended for SARS-CoV-2.

Universal precautions should be routinely followed in dental clinics. They are critical for avoiding the transmission of SARS-CoV-2 virus to children as well as transmission from infected children to healthcare professionals. An infection prevention checklist should be used, including administrative measures, infection prevention education and training, dental healthcare personnel safety, programme evaluation, hand hygiene, personal protective equipment (PPE), respiratory hygiene/cough etiquette, sharps safety, safe injection practices, sterilisation and disinfection of patient-care items and devices, environmental infection prevention and control, and dental unit water quality.^{57,58}

2.3 | Country-specific approaches and recommendations

The WHO has described a pandemic as having six different phases.⁵⁹ Countries will be in different phases at different times; therefore, it is not possible to give

TABLE 1 Summary of reported paediatric cases on COVID-19 (as of 31 March 2020)

Authors	Country	No. of cases reported	Age range
Bi et al ⁹	China	20	-
Cai et al ¹⁰	China	10	3 m to 11 y
CDC ¹¹	USA	123	<19 y
Chan et al ¹²	China	1	<1 y
Chang et al ¹³	China	1	10 y
Chen et al ¹⁴	China	9	<1 y
Chen et al ¹⁵	China	4	<1 y
Choe et al ¹⁶	South Korea	480	0-19 y
Cui et al ¹⁷	China	1	<1 y
D'Antiga ¹⁸	Italy	3	-
Dong et al ⁷	China	731	7 y (median)
Dong et al ¹⁹	China	1	<1 y
Fan et al ²⁰	China	2	<1 y
Henry et al ²¹	International	82	10 y
Ji et al ²²	China	2	>15 y
Kam et al ²³	Singapore	1	<1 y
Li et al ²⁴	China	5	10 m to 6 y
Liu et al ²⁵	China	6	1-7 y
Liu et al ²⁶	China	4	-
Livingston and Bucher ²⁷	Italy	270	<18 y
Lou et al ²⁸	China	3	6 m to 8 y
Lu et al ²⁹	China	171	-
Mizumoto et al ³⁰	Japan	10	0-19 y
Pan et al ³¹	China	1	3 y
Qiu et al ³²	China	36	0-16 y
Tang et al ³³	China	26	1-13 y
Wang et al ³⁴	China	1	<1 y
Wang et al ³⁵	China	1	<1 y
Wang et al ³⁶	China	34	7 y (median)
Wei et al ³⁷	China	9	<1 y
Wu et al ³⁸	China	965	<19 y
Xia et al ³⁹	China	20	-
Xing et al ⁴⁰	China	3	5 and 6 y
Xu et al ⁴¹	China	2	10 to 11 y
Yu et al ⁴²	China	7	<1 y
Zeng et al ⁴³	China	3	<1 y
Zhang et al ⁴⁴	China	34	1 m to 12 y
Zhu et al ⁴⁵	China	10	s<1 y

Abbreviations: CDC, Center for Disease Control and Prevention; m, months; USA, United States of America; y, years.

universal guidelines, so following local updated guidelines

are essential. All the above applies to child patients during the acute phase of COVID-19 pandemic, and the treatment choices and planning may vary during the next phases. Countries have put different measures in place for the overall delivery of dental care even during the 'Widespread Human Infection' phases of the pandemic leading up to the peaks of infection. These vary from those where dental practices remain open but there are screening activities and additional cross-infection measures. These measures generally seem to fit for countries who managed to contain the disease spread quite quickly as seen in Singapore, for example where isolation and testing were put in place within a very short space of the infection being suspected as being present in the country.⁶⁰ At the other end of the spectrum, some countries have closed all dental practices. Some, such as the UK, have all cases triaged by telephone and attending only for very basic treatment in designated centres. In Brazil, The National Health Surveillance Agency (ANVISA) has recommended that only emergency and urgent dental care should be performed (from 20 March 2020), and all private offices have to stop elective treatments. Dental professionals (primary and secondary care) that work for the National Health Service (SUS) have been allocated to help other health professionals in the fast-track for COVID-19.⁶¹ Between these two examples, there are a myriad of service delivery models.

There are a number of country-specific measures currently in place. For example, in the United States (US), Telephone Health (Telehealth) systems have been introduced, and United States Department of Health and Human Services has relaxed the Health Insurance Portability and Accountability Act (HIPAA) regulations in order to enable free and transparent Telehealth services to patients during the COVID-19 public health emergency.⁶² In Brazil, the Ministry of Health has also implemented regulations for Telehealth services to reduce disease transmission.⁶³ Even though countries may be limiting dental care to only emergency provision, the recommendations between countries may differ. There are some differences between recently published documents on urgent non-emergency, and emergency dental procedures by the American Dental Association,⁶⁴ and the UK Scottish Dental Clinical Effectiveness Practice guidelines.⁶⁵ The American Academy of Pediatric Dentistry has also produced an algorithm specific to managing children with emergency dental conditions.^{66,67} As yet, it is difficult to give standard recommendations regarding personal protective equipment (PPE). The use of N-95 respirator masks in the United States and use of filtering facepiece respirator (FFP) masks in Europe are strongly indicated in managing children but this is not a universal recommendation across all countries.⁶⁸ The evidence is lacking for mask use in some areas.⁶⁹ The recommendations provided by

the Center for Disease Control and Prevention (CDC) or other local guidelines that may supersede these should be strictly followed when placing on and removing personal protective equipment used for treating children infected by COVID-19.⁵⁸

2.4 | The role of guidelines and professional judgement

Guidance cannot ever cover all possible circumstances, and professional judgement must be exercised to make decisions around whether or not to provide treatment. Treatment should only be provided when local, regional, and national guidelines are adhered to as far as possible and, in the opinion of the dental professional, is safe for child patients, their accompanying carer, and the dental professional. During this COVID-19 pandemic, universal infection control procedures are of utmost importance with extreme vigilance and championing required by all. Globally, many primary and secondary dental services have been suspended, with many countries providing telephone-based triage systems to identify those patients requiring urgent or emergency intervention.^{59,63} Where dental professionals must offer assessment and/or treatment on a face-to-face basis, they should record all precautions that have been put in place to reduce the risk of cross-infection during treatment and evidence comprehensive risk assessment completion. This includes managing the risk from the treatment itself by carrying out the least invasive treatment possible, and avoiding AGPs.^{57,68}

Healthcare guidance is being updated with alarming frequency, and confusion as to how best proceed in a care setting is both evident and widespread. It is of the highest importance that all members of the dental team acknowledge and act upon their professional responsibility to ensure they are absolutely contemporary in their understanding of current guidance. Additionally, dental teams should be familiar with treatment options that minimise or eliminate AGPs—many of which are founded on contemporary cardiology, well documented in the scientific literature, and minimally invasive by their nature. It must also be appreciated that pandemic experience and staging will differ geographically. Once practice restrictions begin to be eased, continued management of dental disease with contemporary dentistry through minimally interventive concepts and other non-AGPs, whilst viral transmission risk remains high, will be pertinent. These include atraumatic restorative treatment,⁷⁰ sealing in carious lesions using fissure sealants,⁷¹ silver diamine fluoride,⁷² selective caries removal,⁷³ and the Hall Technique.⁷⁴ The importance of toothbrushing with fluoridated toothpaste to prevent tooth decay developing should continue to be emphasised

during contact with patients, and there are opportunities being taken for dentists to carry out telephone and video consultations with parents to promote positive oral health behaviours.

3 | CONCLUSIONS

Although reported clinical manifestations of children's COVID-19 are generally less severe than those of adult patients, young children, and particularly infants, remain vulnerable to infection and pose a significant transmission risk. Dental teams must ensure they remain current in their understanding of local, regional, and national guidance in a climate of uncertainty and frequent change to optimise safety for dental care providers and patients. Dentists who treat children during this pandemic should enact universal infection control procedures to the highest standard and champion this behaviour through their teams. Opportunities to promote preventive dental behaviours should be taken. Contemporary, minimally invasive procedures that minimise or eliminate aerosol generation should be employed where intervention is indicated throughout the pandemic, and in future as and when practice restrictions ease.

DISCLAIMER

The views expressed in this editorial represent the views of the authors and not necessarily those of their host institutions. It has not been peer-reviewed, and it does not replace the clinical judgement of the professional. All sources included should be checked to ensure they are still current.

KEYWORDS

children, coronavirus, COVID-19, dentistry, paediatric dentistry

Sreekanth Kumar Mallineni¹
Nicola P. Innes²
Daniela Procida Raggio³
Mariana Pinheiro Araujo²
Mark D. Robertson²
Jayakumar Jayaraman⁴

¹Pediatric Dentistry, Department of Preventive Dental Science, College of Dentistry, Majmaah University, Al-Zulfi, Saudi Arabia

²Child Dental and Oral Health, School of Dentistry, University of Dundee, Dundee, UK

³Department of Paediatric Dentistry, School of Dentistry, University of Sao Paulo, Sao Paulo, Brazil

⁴Department of Developmental Dentistry, University of Texas Health School of Dentistry, San Antonio, TX, USA

Correspondence

Jayakumar Jayaraman, BDS, MDS, MPaed Dent RCSEd, FDS RCSEd, PhD, Department of Developmental Dentistry, University of Texas Health School of Dentistry, 7703 Floyd Curl Drive, San Antonio, Texas 78229, USA.
Email: jayakumar83@hotmail.com

REFERENCES

1. World Health Organization. [https://www.who.int/news-room/detail/30-01-2020-statement-on-the-second-meeting-of-the-international-health-regulations-\(2005\)-emergency-committee-regarding-the-outbreak-of-novel-coronavirus-\(2019-ncov\)](https://www.who.int/news-room/detail/30-01-2020-statement-on-the-second-meeting-of-the-international-health-regulations-(2005)-emergency-committee-regarding-the-outbreak-of-novel-coronavirus-(2019-ncov)). Accessed March 30, 2020.
2. Coronavirus Resources Center, Johns Hopkins University of Medicine. <https://coronavirus.jhu.edu/map.html>. Accessed March 31, 2020.
3. Meng L, Hua F, Bian Z. Coronavirus disease 2019 (Covid-19): emerging and future challenges for dental and oral medicine. *J Dent Res*;2020. <https://doi.org/10.1177/0022034520914246>
4. Peng X, Xu X, Li Y, Cheng L, Zhou X, Ren B. Transmission routes of 2019-nCoV and controls in dental practice. *Int J Oral Sci*. 2020;12:9.
5. Lauer SA, Grantz KH, Bi Q, et al. The incubation period of coronavirus disease 2019 (COVID-19) from publicly reported confirmed cases: estimation and application. *Annals Int Med*. 2020. <https://doi.org/10.7326/M20-0504>
6. Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan. *China. Lancet*. 2020;395(10223):497-506.
7. Dong Y, Mo X, Hu Y, et al. Epidemiological characteristics of 2143 pediatric patients with 2019 coronavirus disease in China. *Pediatrics*. 2020. <https://doi.org/10.1542/peds.2020-0702>
8. Lu X, Zhang L, Du H, et al. SARS-CoV-2 Infection in Children. *N Engl J Med*. 2020. <https://doi.org/10.1056/NEJMc2005073>
9. Bi Q, Wu Y, Mei S, et al. Epidemiology and Transmission of COVID-19 in Shenzhen China: analysis of 391 cases and 1,286 of their close contacts. *medRxiv*. 2020. <https://doi.org/10.1101/2020.03.03.20028423>
10. Cai J, Xu J, Lin D, et al. A Case Series of children with 2019 novel coronavirus infection: clinical and epidemiological features. *Clin Infect Dis*. 2020. <https://doi.org/10.1093/cid/ciaa198>
11. Bialek S, Boundy E, Bowen V, et al. Severe outcomes among patients with coronavirus disease 2019 (COVID-19) — United States, February 12–March 16, 2020. *MMWR Morb Mortal Wkly Rep*. 2020;69(12):343-346.
12. Chan JF, Yuan S, Kok KH, et al. A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating person-to-person transmission: a study of a family cluster. *Lancet*. 2020;395(10223):514-523.
13. Chang D, Lin M, Wei L, et al. Epidemiologic and clinical characteristics of novel coronavirus infections involving 13 patients outside Wuhan, China. *JAMA*. 2020;323(11):1092.
14. Chen H, Guo J, Wang C, et al. Clinical characteristics and intra-uterine vertical transmission potential of COVID-19 infection in nine pregnant women: a retrospective review of medical records. *Lancet*. 2020;395(10226):809-815.
15. Chen C, Cao M, Peng L, et al. Coronavirus disease-19 among children outside Wuhan, China. *Lancet Child Adolesc Health*. 2020. <http://dx.doi.org/10.2139/ssrn.3546071>

16. Choe YJ. Coronavirus disease-19: The First 7,755 Cases in the Republic of Korea. *medRxiv*. 2020. <https://doi.org/10.1101/2020.03.15.20036368>
17. Cui Y, Tian M, Huang D, et al. A 55-day-old female infant infected with COVID 19: presenting with pneumonia, liver injury, and heart damage. *J Infect Dis*. 2020. <https://doi.org/10.1093/infdis/jiaa113>
18. D'Antiga L. Coronaviruses and immunosuppressed patients. The facts during the third epidemic. *Liver Transpl*. 2020. <https://doi.org/10.1002/lt.25756>
19. Dong L, Tian J, He S, et al. Possible vertical transmission of SARS-CoV-2 from an infected mother to her newborn. *JAMA*. 2020. <https://doi.org/10.1001/jama.2020.4621>
20. Fan C, Lei D, Fang C, et al. Perinatal transmission of COVID-19 associated SARS-CoV- 2: Should we worry? *Clin Infect Dis*. 2020. <https://doi.org/10.1093/cid/ciaa226>
21. Henry BM, Oliveira MHS. Preliminary epidemiological analysis on children and adolescents with novel coronavirus disease 2019 outside Hubei Province, China: an observational study utilizing crowdsourced data. *medRxiv*. 2020. <https://doi.org/10.1101/2020.03.01.20029884>
22. Ji LN, Chao S, Wang YJ, et al. Clinical features of pediatric patients with COVID-19: a report of two family cluster cases. *World J Pediatr*. 2020. <https://doi.org/10.1007/s12519-020-00356-2>. [Epub ahead of print].
23. Kam KQ, Yung CF, Cui L, et al. A well infant with coronavirus disease 2019 (COVID-19) with high viral load. *Clin Infect Dis*. 2020. <https://doi.org/10.1093/cid/ciaa201>
24. Li W, Cui H, Li K, Fang Y, Li S. Chest computed tomography in children with COVID-19 respiratory infection. *Pediatr Radiol*. 2020. <https://doi.org/10.1007/s00247-020-04656-7>
25. Liu H, Liu F, Li J, Zhang T, Wang D, Lan W. Clinical and CT imaging features of the COVID-19 pneumonia: focus on pregnant women and children. *J Infect*. 2020. <https://doi.org/10.1016/j.jinf.2020.03.007>
26. Liu W, Zhang Q, Chen J, et al. Detection of Covid-19 in children in early January 2020 in Wuhan, China. *N Engl J Med*. 2020. <https://doi.org/10.1056/NEJMc2003717>
27. Livingston E, Bucher K. Coronavirus Disease 2019 (COVID-19) in Italy. *JAMA*. 2020. <https://doi.org/10.1001/jama.2020.4344>
28. Lou XX, Shi CX, Zhou CC, Tian YS. Three children who recovered from novel coronavirus 2019 pneumonia. *J Paediatr Child Health*. 2020. <https://doi.org/10.1111/jpc.14871>
29. Lu X, Zhang L, Du H, et al. SARS-CoV-2 Infection in Children. *N Engl J Med*. 2020. <https://doi.org/10.1056/NEJMc2005073>
30. Mizumoto K, Omori R, Nishiura H. Age specificity of cases and attack rate of novel 2 coronavirus disease (COVID-19). *medRxiv*. 2020. <https://doi.org/10.1101/2020.03.09.20033142>
31. Pan X, Chen D, Xia Y, et al. Asymptomatic cases in a family cluster with SARS-CoV-2 infection. *Lancet Infect Dis*. 2020. [https://doi.org/10.1016/S1473-3099\(20\)30114-6](https://doi.org/10.1016/S1473-3099(20)30114-6)
32. Qiu H, Wu J, Hong L, Luo Y, Song Q, Chen D. Clinical and epidemiological features of 36 children with coronavirus disease 2019 (COVID-19) in Zhejiang, China: an observational cohort study. *Lancet Infect Dis*. 2020. [https://doi.org/10.1016/S1473-3099\(20\)30198-5](https://doi.org/10.1016/S1473-3099(20)30198-5)
33. Tang A, Xu W, Shen M, et al. A retrospective study of the clinical characteristics of COVID-19 infection in 26 children. *medRxiv*. 2020. <https://doi.org/10.1101/2020.03.08.20029710>
34. Wang S, Guo L, Chen L, et al. A case report of neonatal COVID-19 infection in China. *Clin Infect Dis*. 2020. <https://doi.org/10.1093/cid/ciaa225>
35. Wang X, Zhou Z, Zhang J, Zhu F, Tang Y, Shen X. A case of 2019 Novel Coronavirus in a pregnant woman with preterm delivery. *Clin Infect Dis*. 2020. <https://doi.org/10.1093/cid/ciaa200>
36. Wang XF, Yuan J, Zheng YJ, et al. Clinical and epidemiological characteristics of 34 children with 2019 novel coronavirus infection in Shenzhen. *Zhonghua Er Ke Za Zhi*. 2020;58:E008.
37. Wei M, Yuan J, Liu Y, Fu T, Yu X, Zhang ZJ. Novel coronavirus infection in hospitalized infants under 1 year of age in China. *JAMA*. 2020. <https://doi.org/10.1001/jama.2020.2131>
38. Wu Z, McGoogan JM. Characteristics of and important lessons from the coronavirus disease 2019 (COVID-19) outbreak in China. *JAMA*. 2020. <https://doi.org/10.1001/jama.2020.2648>
39. Xia W, Shao J, Guo Y, Peng X, Li Z, Hu D. Clinical and CT features in paediatric patients with COVID-19 infection: different points from adults. *Pediatr Pulmonol*. 2020. <https://doi.org/10.1002/ppul.24718>
40. Xing Y, Ni W, Wu Q, et al. Prolonged presence of SARS-CoV-2 in feces of paediatric patients during the convalescent phase. *medRxiv*. 2020. <https://doi.org/10.1101/2020.03.11.20033159>
41. Xu XW, Wu XX, Jiang XG, Xu KJ, Ying LJ, Ma CL. Clinical findings in a group of patients infected with the 2019 novel coronavirus (SARS-Cov-2) outside of Wuhan, China: retrospective case series. *BMJ*. 2020. <https://doi.org/10.1136/bmj.m606>
42. Yu N, Li W, Kang Q, et al. Clinical features and obstetric and neonatal outcomes of pregnant patients with COVID-19 in Wuhan, China: a retrospective, single-centre, descriptive study. *Lancet Infect Dis*. 2020. [https://doi.org/10.1016/S1473-3099\(20\)30176-6](https://doi.org/10.1016/S1473-3099(20)30176-6)
43. Zeng L, Xia S, Yuan W, et al. Neonatal Early-Onset Infection With SARS-CoV-2 in 33 Neonates Born to Mothers With COVID-19 in Wuhan, China. *JAMA Pediatr*. 2020. <https://doi.org/10.1001/jamapediatrics.2020.0878>
44. Zhang SC, Gu J, Chen Q, Deng N. Clinical characteristics of 34 children with coronavirus disease-2019 in the West of China: a multiple-center case. *medRxiv*. 2020. <https://doi.org/10.1101/2020.03.12.20034686>
45. Zhu H, Wang L, Fang C, et al. Clinical analysis of 10 neonates born to mothers with 2019-nCoV pneumonia. *Transl Pediatr*. 2020;9(1):51-60.
46. Sun P, Lu X, Xu C, Sun W, Pan B. Understanding of COVID 19 based on current evidence. *J Med Virol*. 2020. <https://doi.org/10.1002/jmv.25722>
47. Illinois Department of Public Health. <http://dph.illinois.gov/news/public-health-officials-announce-first-death-infant-coronavirus-disease>. Accessed March 29, 2020.
48. National Broadcasting Corporation, New York. <https://www.nbcnewyork.com/news/local/nyc-virus-deaths-leap-from-0-to-776-in-15-days-emergency-hospital-help-arrives-monday/2350357/>. Accessed March 30, 2020.
49. The Daily Mail. <https://www.dailymail.co.uk/news/article-8171301/12-year-old-infected-coronavirus-dies-Belgium.html>. Accessed March 31, 2020.
50. The Daily Mail. <https://www.dailymail.co.uk/news/article-8173253/Boy-13-dies-testing-positive-coronavirus-London-hospital-fundraiser-says.html>. Accessed March 31, 2020.
51. The Sun. <https://www.thesun.co.uk/news/11268073/french-girk-youngest-coronavirus-victim-cough/>. Accessed March 31, 2020.
52. Teenager's Death in California is linked to Coronavirus. The New York Times. <https://www.nytimes.com/2020/03/24/us/california-coronavirus-death-child.html>. Accessed March 30, 2020.

53. To KK, Tsang OT, Yip CC, et al. Consistent detection of 2019 novel coronavirus in saliva. *Clin Infect Dis*. 2020. <https://doi.org/10.1093/cid/ciaa149>
54. Fan C, Lei D, Fang C, et al. Perinatal transmission of COVID-19 associated SARS-CoV- 2: Should we worry? *Clin Infect Dis*. 2020. <https://doi.org/10.1093/cid/ciaa226>
55. Tran K, Cimon K, Severn M, Pessoa-Silva CL, Conly J. Aerosol generating procedures and risk of transmission of acute respiratory infections to healthcare workers: a systematic review. *PLoS ONE*. 2012;7:e35797.
56. van Doremalen N, Bushmaker T, Morris DH, et al. Aerosol and surface stability of SARS-CoV-2 as compared with SARS-CoV-1. *New Eng J Med*. 2020. <https://doi.org/10.1056/NEJMc2004973>
57. Smales FC, Samaranyake LP. Maintaining dental education and specialist dental care during an outbreak of a new coronavirus infection. Part 2: Control of the disease, then elimination. *Br Dent J*. 2003;195:679-681.
58. Personal Protective Equipment. Center for Disease Control and Prevention. <https://www.cdc.gov/hai/pdfs/ppe/ppe-sequence.pdf>. Accessed March 30, 2020.
59. World Health Organization Pandemic Phase Descriptions. https://www.who.int/influenza/resources/documents/pandemic_phase_descriptions_and_actions.pdf. Accessed March 31, 2020.
60. Dental Council of Singapore. <https://www.healthprofessionals.gov.sg/sdc>. Accessed March 30, 2020.
61. Brazil National Health System (Sistema Unico de Saude). Fast-track for primary care in places with community transmission.. http://maismedicos.gov.br/images/fluxo_bolso_17mar20_3.pdf. Accessed March 31, 2020.
62. Notification of Enforcement Discretion for Telehealth Remote Communications During the COVID-19 Nationwide Public Health Emergency. <https://www.hhs.gov/hipaa/for-professionals/special-topics/emergency-preparedness/notification-enforcement-discretion-telehealth/index.html>. Accessed March 31, 2020.
63. Brazil National Health System (Sistema Unico de Saude). Dental care during COVID-19. http://website.cfo.org.br/wp-content/uploads/2020/03/COVID-19_ATENDIMENTO-ODONTOLOGICO-NO-SUS.pdf. Accessed March 31, 2020.
64. American Dental Association. What constitutes a medical emergency? https://success.ada.org/~media/CPS/Files/Open%20Files/ADA_COVID19_Dental_Emergency_DDS.pdf?_ga=2.236419265.19784197.1584891239-1970975310.1584891239. Accessed March 30, 2020.
65. Scottish Dental Clinical Effectiveness Program. <http://www.sdcep.org.uk/wp-content/uploads/2020/03/SDCEP-MADP-COVID-19-guide-300320.pdf>. Accessed March 31, 2020.
66. American Academy of Pediatric Dentistry. <https://www.aapd.org/about/about-aapd/news-room/covid-19/>. Accessed March 30, 2020.
67. Meyer BD, Casamassimo P, Vann WF Jr. An algorithm for managing emergent dental conditions for children. *J Clin Pediatr Dent*. 2019;43:201-206.
68. Smales FC, Samaranyake LP. Maintaining dental education and specialist dental care during an outbreak of a new coronavirus infection. Part 1: a deadly viral epidemic begins. *Br Dent J*. 2003;195:557-561.
69. Centre for Evidence Based Medicine, University of Oxford. <https://www.cebm.net/covid-19/what-is-the-efficacy-of-standard-face-masks-compared-to-respirator-masks-in-preventing-covid-type-respiratory-illnesses-in-primary-care-staff/>. Accessed March 30, 2020.
70. De Amorim RG, Frencken JE, Raggio DP, Chen X, Hu X, Leal SC. Survival percentages of atraumatic restorative treatment (ART) restorations and sealants in posterior teeth: an updated systematic review and meta-analysis. *Clinical Oral Invest*. 2018;22:2703-2725.
71. Schwendicke F, Jäger AM, Paris S, Hsu LY, Tu YK. Treating pit-and-fissure caries: a systematic review and network meta-analysis. *J Dent Res*. 2015;94:522-533.
72. Seifo N, Cassie H, Radford JR, Innes NP. Silver diamine fluoride for managing carious lesions: an umbrella review. *BMC Oral Health*. 2019;19:145.
73. Li T, Zhai X, Song F, Zhu H. Selective versus non-selective removal for dental caries: a systematic review and meta-analysis. *Acta Odontol Scand*. 2018;76:135-140.
74. Innes NP, Evans DJ, Stirrups DR. Sealing caries in primary molars: randomized control trial, 5-year results. *J Dent Res*. 2011;90:1405-1410.

How to cite this article: Mallineni SK, Innes NP, Raggio DP, Araujo MP, Robertson MD, Jayaraman J. Coronavirus disease (COVID-19): Characteristics in children and considerations for dentists providing their care. *Int J Paediatr Dent*. 2020;00:1–6. <https://doi.org/10.1111/ipd.12653>